

Dosing Information for Healthcare Professionals

Infants' **TYLENOL** Children's **TYLENOL**

Use this chart to determine the proper dose of **TYLENOL**® for your patient. If possible, use weight to dose; otherwise use age.

**DOSE: Every 4 hours as needed.
Do not give more than 5 doses in 24 hours.**

WEIGHT	AGE	Infants' TYLENOL ® Oral Suspension Active Ingredient: Acetaminophen 160 mg (in each 5 mL)	Children's TYLENOL ® Oral Suspension Active Ingredient: Acetaminophen 160 mg (in each 5 mL or 1 tsp)
6-11 lbs	0-3 mos	1.25 mL	—
12-17 lbs	4-11 mos	2.5 mL	—
18-23 lbs	12-23 mos	3.75 mL	—
24-35 lbs	2-3 yrs	5 mL	5 mL (1 tsp)
36-47 lbs	4-5 yrs	—	7.5 mL (1½ tsp)
48-59 lbs	6-8 yrs	—	10 mL (2 tsp)
60-71 lbs	9-10 yrs	—	12.5 mL (2½ tsp)
72-95 lbs	11 yrs	—	15 mL (3 tsp)

mL = milliliter tsp = teaspoonful

Remind parents and caregivers to:

- Read and follow the label on all **TYLENOL**® products
- Take every 4 hours as needed.
- Do NOT exceed more than 5 doses in 24 hours
- Do NOT use with any other product containing acetaminophen
- Use only the dosing device that comes with a specific product

All Infants' **TYLENOL**® and Children's **TYLENOL**® Oral Suspension products have the same acetaminophen concentration (160 mg/5 mL)

Dosing Information for Healthcare Professionals

Children's **Motrin** Concentrated **Motrin** Infants' Drops

Use this chart to determine the proper dose of **MOTRIN**® for your patient. If possible, use weight to dose; otherwise use age.

**DOSE: Every 6-8 hours as needed.
Do not give more than 4 doses in 24 hours.**

WEIGHT	AGE	Concentrated MOTRIN ® Infants' Drops Active Ingredient: Ibuprofen 50 mg (NSAID)* (in each 1.25 mL)	Children's MOTRIN ® Oral Suspension Active Ingredient: Ibuprofen 100 mg (NSAID)* (in each 5 mL or 1 tsp)
6-11 lbs	0-5 mos	—	—
12-17 lbs	6-11 mos	1.25 mL	—
18-23 lbs	12-23 mos	1.875 mL	—
24-35 lbs	2-3 yrs	—	5 mL (1 tsp)
36-47 lbs	4-5 yrs	—	7.5 mL (1½ tsp)
48-59 lbs	6-8 yrs	—	10 mL (2 tsp)
60-71 lbs	9-10 yrs	—	12.5 mL (2½ tsp)
72-95 lbs	11 yrs	—	15 mL (3 tsp)

mL = milliliter tsp = teaspoonful

Remind parents and caregivers to:

- Read and follow the label on all **MOTRIN**® products
- Take every 6-8 hours as needed. Do NOT exceed more than 4 doses in 24 hours
- Do NOT administer longer than 10 days, unless directed by a doctor
- Ask a doctor or pharmacist before use if the child is taking any other drug containing an NSAID* (prescription or non-prescription)
— **MOTRIN**® contains ibuprofen
- Use only the dosing device that comes with a specific product

*Nonsteroidal anti-inflammatory drug
©McNEIL-PPC, Inc. 2013 • CTCM-0144

My healthcare professional recommends...


Children's Advil

Relief you can trust

Children's Advil®

Lasts Up To
8
Hours


Find the right dose on the chart below. If possible, use weight to dose otherwise use age.



Infants' Advil®
Concentrated Drops
for ages 6-23 months

ACTIVE INGREDIENT
Ibuprofen 50 mg (NSAID)[†]
in each 1.25 mL


Fever reducer; relieves minor aches and pains due to the common cold, flu, headaches, and toothaches



Children's Advil®
Suspension
for ages 2-11 years

ACTIVE INGREDIENT
Ibuprofen 100 mg (NSAID)[†]
in each 5 mL

Fever reducer; relieves minor aches and pains due to the common cold, flu, sore throat headaches, and toothaches



Junior Strength Advil®
Chewables
for ages 6-11 years

ACTIVE INGREDIENT
Ibuprofen 100 mg (NSAID)[†]
in each tablet

Fever reducer; relieves minor aches and pains due to the common cold, flu, sore throat headaches, and toothaches



Junior Strength Advil®
Tablets
for ages 6-11 years

ACTIVE INGREDIENT
Ibuprofen 100 mg (NSAID)[†]
in each tablet

Fever reducer; relieves minor aches and pains due to the common cold, flu, sore throat headaches, and toothaches

		GIVE YOUR CHILD THE DOSE INDICATED BELOW EVERY 6-8 HOURS, NOT TO EXCEED 4 DOSES/24 HOURS					
PATIENT DOSING	Weight	Age					
		under 6 months	Ask a doctor				
	12-17 pounds	6-11 months	1.25 mL [‡]	Ask a doctor about using Infants' Advil®			
		12-23 months	1.875 mL [‡]				
	18-23 pounds	2-3 years	Ask a doctor about using Children's or Junior Strength Advil®	1 tsp [‡]		Ask a doctor about using Infants' or Children's Advil®	
		4-5 years		1½ tsp [‡]			
		6-8 years		2 tsp [‡]	2 tablets		2 tablets
		9-10 years		2½ tsp [‡]	2½ tablets		2 tablets
11 years		3 tsp [‡]		3 tablets	3 tablets		

For 12+ years, use adult strength Advil®